


Hip Resurfacing/Total Hip Arthroplasty Patients

Phase One *10% Weight Bearing* *Slow Program*

revised 08/08/2024 tpg/law

1. Points to remember for 6 weeks:

- Avoid flexing (bending) your hip greater than 90 degrees. Before standing up, scoot to the edge and slide your operated leg forward. Before sitting down, slide your operated leg forward, sit on the edge and then scoot back. When getting in and out of the car have the seat slid all the way back and the seat back reclined. Lean back and have your shoulders behind your hips as you bring your legs in and out of the car. When you are in the car you can have the seat positioned for comfort.
- Avoid hip internal rotation (turning your leg inward).
- Do not do straight leg raises for exercise. Okay to lift your leg to move it on and off the bed and in and out of the car when comfortable.
- Do not cross your legs when your hips and knees are flexed (bent). Okay to cross at your ankles.
- Okay to lie on either side. Okay to use a pillow(s) between your legs for comfort when on your side.
- Okay to reach towards your foot when sitting to put on socks and shoes if your knee can fall out to the side (hip external rotation). It may be most comfortable to rest your foot on a foot stool, let your knee roll out and then bend forward. Your foot must be on the floor or supported on a foot stool.



2. Slow Walking Program:

1. Two crutches or a walker at all times 10% weight bearing for four weeks.
****When you are standing you can have equal weight on your legs.****
2. Then two crutches or a walker at all times weight bearing as tolerated for two weeks.
3. Then one crutch or a cane held in the opposite hand for one month.
****Functional/necessity walking only for six months when fracture risk has passed.****

Crutch/Walker Walking Sequence

Level Surface- Move your crutches or walker forward first then your operated leg then your non-operated leg.

Up Steps- “up with the strong”. One step at a time. Your non-operated leg will go up first then your operated leg then the crutches. You can use a handrail on both sides or a handrail on one side and crutch or cane on the other side. You cannot use a walker on multiple steps.


Down Steps- “down with the weak”. One step at a time. Your crutches will go down first then your operated leg then your non-operated leg. You can use a handrail on both sides or a handrail on one side and crutch or cane on the other side. You cannot use a walker on multiple steps.

****One step at a time on stairs for 10 weeks****

3. Perform the following exercises:


a. Ankle Pumps

Sitting or lying, bend your feet up and down at your ankles. Move through the full range of motion for blood circulation. Do this exercise 30 times every waking hour for the first two weeks.



b. Isometrics

Sitting or lying with your legs straight, tighten the muscles on the front of your thighs (quadriceps) by pressing your knees flat and simultaneously squeeze your buttocks muscles. Hold five seconds. Relax. Repeat. Start with 30 times two times per day. Okay to do more but don't make yourself sore. Do not hold your breath during this exercise.




c. Heel Slides

Lying flat on your bed with your legs straight, gently slide your heel towards your buttocks. Then slide your heel out until your knee is straight. Relax. Repeat. Do this exercise 10-15 times two times per day only. You have to do this on your operated side. Optional on the other side. Do not do both legs at the same time.



d. Short Arc Quads

Lying flat on your bed with your knee resting on a rolled towel 3-4 inches wide, lift your heel off your bed straightening your knee. Keep your knee on the roll. Hold five seconds. Slowly lower. Relax. Repeat. Do this exercise 15-30 times two times per day only. You have to do this on your operated side. Optional on the other leg. Do not do both legs at the same time.



You may not use a stationary bike or elliptical for 10 weeks on the slow program. You may not use a stair stepper or rowing machine or do mountain climbers for six months. At week three you can start arm exercises. You must be seated and can use free weights, resistance bands or machines. Any weight is OK as long as you don't push with your legs or stabilize with your core. Mini-crunches lying in your bed and pull-ups are OK at week three but no pushups, no sit-ups, no bridges, no planks for six weeks.